



Metformin vildagliptin tablets (II)

[Indications] Currently no data available

[Usage and Dosage] When using this product for hypoglycemic treatment, the dosage should be customized according to the patient's current treatment plan, efficacy, and tolerance to the drug. But the maximum daily dose of Vigagliptin should not exceed the recommended 100mg. The usual dosing regimen is twice a day, one tablet in the morning and one tablet in the evening. Taking this product during or after meals can alleviate gastrointestinal symptoms of metformin (see pharmacokinetics). For patients with poor blood glucose control despite reaching the maximum tolerable dose of metformin monotherapy, the starting dose of Benchang is equivalent to 50mg of Veragliptin twice daily (total daily dose 100mg), plus the dose of metformin currently being taken. (See instructions for details)

[Adverse reactions] There is currently no clinical efficacy trial data for this product, but trials have shown that it is bioequivalent with the combined use of Vigagliptin and Metformin. The following data comes from a study of the combined use of Vigagliptin and Metformin tablets, in which Vigagliptin tablets were used as an additional therapeutic drug to

Metformin tablets. There is currently no research on the addition of metformin in the treatment of Vigagliptin.

【 Taboos 】 It is prohibited for those who are known to be allergic to any of the ingredients in this product, such as Vigagliptin, Metformin.

[Note] For patients who require insulin treatment, this product cannot replace insulin. This product is not suitable for patients with type 1 diabetes. Please read the instructions carefully and follow the doctor's instructions when using.